=> d his

(FILE 'HOME' ENTERED AT 13:14:38 ON 24 JAN 2011)

FILE 'REGISTRY' ENTERED AT 13:15:03 ON 24 JAN 2011

L1 STRUCTURE UPLOADED

L2 2 S L1

L3 24 S L1 SSS FUL

FILE 'CAPLUS' ENTERED AT 13:15:54 ON 24 JAN 2011

L4 6 S L3

=> d ibib abs hitstr total

L4 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2011 ACS on STN

ACCESSION NUMBER: 2007:1440598 CAPLUS

DOCUMENT NUMBER: 148:229130

TITLE: 2,3-Benzodiazepine-type AMPA receptor antagonists and

their neuroprotective effects

AUTHOR(S): Szenasi, Gabor; Vegh, Miklos; Szabo, Geza; Kertesz,

Szabolcs; Kapus, Gabor; Albert, Mihaly; Greff, Zoltan;

Ling, Istvan; Barkoczy, Jozsef; Simig, Gyula;

Spedding, Michael; Harsing, Laszlo G.

CORPORATE SOURCE: Division of Preclinical Research, EGIS Pharmaceuticals

Plc, Budapest, 1165, Hung.

SOURCE: Neurochemistry International (2008), 52(1-2), 166-183

CODEN: NEUIDS; ISSN: 0197-0186

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal LANGUAGE: English

AMPA receptors are fast ligand-gated members of glutamate receptors in neuronal and many types of non-neuronal cells. The heterotetramer complexes are assembled from four subunits (GluR1-4) in region-, development- and function-selective patterns. Each subunit contains three extracellular domains (a large amino terminal domain, an agonist-binding domain and a transducer domain), and three transmembrane segments with a loop (pore forming domain), as well as the intracellular carboxy terminal tail (traffic and conductance regulatory domain). The binding of the agonist (excitatory amino acids and their derivs.) initiates conformational realignments, which transmit to the transducer domain and membrane spanning segments to gate the channel permeable to Na+, K+ and more or less to Ca2+. Several 2,3-benzodiazepines act as non-competitive antagonists of the AMPA receptor (termed also neg. allosteric modulators), which are thought to bind to the transducer domains and inhibit channel gating. Analyzing their effects in vitro, it has been possible to recognize a structure-activity relationship, and to describe the critical parts of the mols. involved in their action at AMPA receptors. Blockade of AMPA receptors can protect the brain from apoptotic and necrotic cell death by preventing neuronal excitotoxicity during pathophysiol. activation of glutamatergic neurons. Animal expts. provided evidence for the potential usefulness of non-competitive AMPA antagonists in the treatment of human ischemic and neurodegenerative disorders including stroke, multiple sclerosis, Parkinson's disease, periventricular leukomalacia and motoneuron disease. 2,3-Benzodiazepine AMPA antagonists can protect against seizures, decrease levodopa-induced dyskinesia in animal models of Parkinson's disease demonstrating their utility for the treatment of a variety of CNS disorders.

IT 439143-67-6 439143-70-1

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(2,3-Benzodiazepine-type AMPA receptor antagonists and their neuroprotective effects)

RN 439143-67-6 CAPLUS

CN Ethanone, 1-[1-(4-amino-3-methylphenyl)-8-chloro-4,5-dihydro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

10/567,598

RN 439143-70-1 CAPLUS

CN Ethanone, 1-[1-(4-amino-3-methylphenyl)-7,8-dichloro-4,5-dihydro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

OS.CITING REF COUNT: 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

REFERENCE COUNT: 164 THERE ARE 164 CITED REFERENCES AVAILABLE FOR

THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2011 ACS on STN

ACCESSION NUMBER: 2007:755669 CAPLUS

DOCUMENT NUMBER: 147:166354

TITLE: Process for preparation of chiral

dihydro-2,3-benzodiazepines as AMPA receptor

antagonists

INVENTOR(S): Ling, Istvan; Barkoczy, Jozsef; Greff, Zoltan;

Szenasi, Gabor; Gigler, Gabor; Kertesz, Szabolcs; Szuecs, Gyula; Albert, Mihaly; Kapus, Gabor; Szabo, Geza; Vegh, Miklos; Agoston, Marta; Levay, Gyoergy;

Moricz, Krisztina; Harsing, Laszlo Gabor

PATENT ASSIGNEE(S): Egis Gyogyszergyar Nyrt., Hung.

SOURCE: PCT Int. Appl., 144pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PA'	TENT	ΝΟ.			KIND DATE						LICAT	DATE						
WO	2007	0774	69		A1	2007	0712			2006-		20061229						
	W:	ΑE,	AG,	AL,	AM,	ΑT,	AU,	AZ,	BA,	BB	, BG,	BR,	BW,	BY,	BZ,	CA,	CH,	
		CN,	CO,	CR,	CU,	CZ,	DE,	DK,	DM,	DZ	, EC,	EE,	EG,	ES,	FΙ,	GB,	GD,	
		GE,	GH,	GM,	GT,	HN,	HR,	HU,	ID,	ΙL	, IN,	IS,	JP,	ΚE,	KG,	KM,	KN,	
		KP,	KR,	KΖ,	LA,	LC,	LK,	LR,	LS,	LΤ	, LU,	LV,	LY,	MA,	MD,	MG,	MK,	
		MN,	MW,	MX,	MY,	MZ,	NA,	NG,	ΝI,	NO	, NZ,	OM,	PG,	PH,	PL,	PT,	RO,	
		RS,	RU,	SC,	SD,	SE,	SG,	SK,	SL,	SM	, SV,	SY,	ТJ,	TM,	TN,	TR,	TT,	
		TZ,	UA,	UG,	US,	UZ,	VC,	VN,	ZA,	ZM	, ZW							
	RW:	ΑT,	BE,	ΒG,	CH,	CY,	CZ,	DE,	DK,	EE	, ES,	FI,	FR,	GB,	GR,	HU,	IE,	
		IS,	ΙT,	LT,	LU,	LV,	MC,	NL,	PL,	PT	, RO,	SE,	SI,	SK,	TR,	BF,	ВJ,	
		CF,	CG,	CI,	CM,	GΑ,	GN,	GQ,	GW,	ML	, MR,	NE,	SN,	TD,	ΤG,	BW,	GH,	
		GM,	KΕ,	LS,	MW,	${ m MZ}$,	NA,	SD,	SL,	SZ	, TZ,	UG,	ZM,	ZW,	ΑM,	ΑZ,	BY,	
		,			RU,	,												
	U 2005001211								HU 2005-1211						2	0051	230	
	2005						2007											
	2005								HU 2005-1212						2	0051	230	
	2005						2008						00061666					
	2006											20061229						
	2633									-			20061229					
EP	1979										2006-							
	R:	,	,	,	,	,		,	,		, ES,		,	,	•	•	,	
						LU,	L∨,	MC,	ΝL,	ΡЬ	, PT,	RO,	SE,	SI,	SK,	TR,	AL,	
TD	2000		HR,				2000	0611		TD	0000	F 400	2.2		0	0061	000	
	2009										2008-				_	0061		
	CN 101389600								CN 2006-80053005 US 2008-159251									
	US 20090233913 ORITY APPLN. INFO.:						2009	091/										
KIOKIT	I APP	υΝ.	TMFO	.:							2005-							
											2005- 2006-					0051		
										WU	2006-	потз	U		vv Z	υυφΙ	449	

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT OTHER SOURCE(S): CASREACT 147:166354; MARPAT 147:166354

GΙ

AB This invention provides four processes for the preparation of chiral dihydro-2,3-benzodiazepines I [wherein X = halo or alkoxy; Y = halo; or X and Y = methylenedioxy; R = alkyl.] or pharmaceutically acceptable salts thereof as AMPA receptor antagonists. For example, II was prepared via several stereoselective approaches. In spreading depression test, II showed antagonistic activity with EC50 of 1.8 \pm 0.1 μ M against AMPA receptor. The compds. are useful neuroprotective agents for the treatment of stoke, epilepsy, schizophrenia, etc. (no data).

IT 943964-01-0P 943964-02-1P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU
 (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES
 (Uses)

(drug candidate; preparation of chiral dihydro-2,3-benzodiazepines as AMPA receptor antagonists)

RN 943964-01-0 CAPLUS

CN Ethanone, 1-[(4R)-1-(4-amino-3-methylphenyl)-8-chloro-4,5-dihydro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

RN 943964-02-1 CAPLUS

CN Ethanone, 1-[(4S)-1-(4-amino-3-methylphenyl)-8-chloro-4,5-dihydro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).

IT 943964-14-5P 943964-18-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(intermediate; preparation of chiral dihydro-2,3-benzodiazepines as AMPA receptor antagonists)

RN 943964-14-5 CAPLUS

CN Ethanone, 1-[(4R)-8-chloro-4,5-dihydro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).

RN 943964-18-9 CAPLUS

CN Ethanone, 1-[(4S)-8-chloro-4,5-dihydro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

6

REFERENCE COUNT:

THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2011 ACS on STN

ACCESSION NUMBER: 2007:389016 CAPLUS

DOCUMENT NUMBER: 147:469380

TITLE: New substituted 2,3-benzodiazepine derivatives, their

use and pharmaceutical compositions containing them INVENTOR(S): Solyom, Sandor; Abraham, Gizella; Berzsenyi, Pal; Andrasi, Ferenc; Szabo, Hilda; Csuzdi, Emese; Hamori,

Tamas; Kertesz, Mariusz; Csillikne, Perczel Viola; Horvath, Gyula; Kurucz, Istvan; Pallagi, Istvan; Toth,

Szilveszter; Toeroek, Katalin; Ling, Istvan

PATENT ASSIGNEE(S): Ivax Gyogyszerkutato Intezet Kft., Hung.

SOURCE: Hung. Pat. Appl., 96pp.

CODEN: HUXXCV

Ι

DOCUMENT TYPE: Patent LANGUAGE: Hungarian

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE		
HU 2004000338	A2	20060928	HU 2004-338	20040203		
PRIORITY APPLN. INFO.:			HU 2004-338	20040203		
GI						

AB The invention concerns the general formula 2,3-benzodiazepines I [R1, R2 = H, C1-3-alkyl; R3 = (un)substituted C5-6-aromatic, saturated or partially saturated

heterocycle containing at least 2 heteroatoms, in which, as heteroatom, there may be oxygen, sulfur or nitrogen atom and in case the heterocycle contains 2 heteroatoms, one of the heteroatoms must be something other than nitrogen atom; R4, R5, R6, R7, R8 = H, C1-3-alkyl, halogen, NO2, NH2, NH(C1-3-alkyl), N(C1-3-alkyl)2, C2-5-acyl, (C2-5-alkoxy)carbonyl, C2-5-alkylaminocarbonyl; R9 = C1-3-alkoxy, halogen; R10 = H, halogen; R9R10 = C1-3-alkylenedioxy] and their isomers and acid addition salts. Thus, $(\pm)-5-(4-aminophenyl)-8-methyl-7-(2-thiazolyl)-8,9-dihydro-7H-1,3-dioxolo[4,5-h][2,3]benzodiazepine I [R1 = R4 = R5 = R7 = R8 = H, R2 = Me, R3 = thiazol-2-yl, R6 = NH2, R9R10 = OCH20] was prepared from <math>(\pm)-8-Methyl-5-(4-nitrophenyl)-8,9-dihydro-7H-1,3-dioxolo[4,5-h][2,3]benzodiazepine (II) via thiocarbamylation with potassium thiocyanate in AcOH, cyclocondensation with BrCH2CH(OEt)2 in DMF, and reduction with H2NNH2·H2O in MeOH/CH2C12 containing catalytic RaNi. The$

CN

invention includes the pharmaceutical compns. that contain the above compds. I, the application of the compds. and the production of pharmaceutical products to treat neurodegenerative diseases. The pharmacol. activity of I was determined

IT 952603-81-5P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and cyclocondensation reactions of; new substituted 2,3-benzodiazepine derivs., their use and pharmaceutical compns. containing them)

RN 952603-81-5 CAPLUS

3H-2,3-Benzodiazepine-3-carbothioamide, 7,8-dichloro-4,5-dihydro-4-methyl-1-(3-methyl-4-nitrophenyl)- (CA INDEX NAME)

$$\begin{array}{c|c} \text{Me} & \text{C1} \\ \text{H}_2\text{N}-\text{C} & \text{N} \\ \text{S} & \text{N} \end{array}$$

ANSWER 4 OF 6 CAPLUS COPYRIGHT 2011 ACS on STN T.4

2007:119526 CAPLUS ACCESSION NUMBER:

DOCUMENT NUMBER: 146:206341

TITLE: Novel substituted 2,3-benzodiazepine derivatives as

AMPA antagonists and their preparation, pharmaceutical compositions, and use in the treatment of diseases

INVENTOR(S): Solyom, Sandor; Abraham, Gizella; Hamori, Tamas; Berzsenyl, Pal; Andrasi, Fenrec; Kurucz, Istvan

PATENT ASSIGNEE(S): IVAX Drug Research Institute, Ltd., Hung.

SOURCE: U.S. Pat. Appl. Publ., 40 pp., Cont.-in-part of U.S.

Ser. No. 358,053.

CODEN: USXXCO

DOCUMENT TYPE: Patent English LANGUAGE:

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070027143	A1	20070201	US 2004-771847	20040203
US 20040152693	A1	20040805	US 2003-358053	20030204
US 6858605	B2	20050222		
PRIORITY APPLN. INFO.:			US 2003-358053 A2	2 20030204
ASSIGNMENT HISTORY FOR U	JS PATEN	T AVAILABLE	IN LSUS DISPLAY FORMAT	
OTHER SOURCE(S):	CASREA	CT 146:20634	11: MARPAT 146:206341	

GΙ

The invention relates to 2,3-benzodiazepine derivs. of formula I, isomers AB and acid addition salts thereof and to pharmaceutical compns. containing the same, as well as to pharmaceutical compns. and methods of using the same suitable for treating conditions associated with muscle spasms, epilepsy, acute and chronic forms of neurodegenerative diseases as well as preventing, treating or alleviating the symptoms of acute and chronic inflammatory disorders. Compds. of formula I wherein, R3 is (un) substituted (un) saturated 5- to 6-membered (hetero) aryl; R4 - R7 is H, halo, C1-3 alkyl, NO2, NH2 and derivs., etc.; R9 is C1-3 alkoxy, and halo; R10 is H and halo; R9R10 together is C1-3 alkylenedioxy; and their stereoisomers and acid-addition salts thereof, are claimed. Example compound II was prepared by cyclization of $(\pm)-8-methyl-5-(4-nitrophenyl)-7$ thiocarbamoyl-8,9-dihydro-7H-1,3-dioxolo[4,5-h][2,3]benzodiazepine with bromoacetaldehyde di-Et acetal. All the invention compds. were evaluated

CN

for their AMPA antagonistic activity (data given).

IT 923271-76-5P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; preparation of substituted 2,3-benzodiazepine derivs. as AMPA antagonists and their use in the treatment of diseases)

RN 923271-76-5 CAPLUS

3H-2,3-Benzodiazepine, 8-chloro-4,5-dihydro-4-methyl-1-(3-methyl-4-nitrophenyl)-3-(5-methyl-1,3,4-thiadiazol-2-yl)- (CA INDEX NAME)

L4 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2011 ACS on STN

ACCESSION NUMBER: 2005:120900 CAPLUS

DOCUMENT NUMBER: 142:219316

TITLE: Process for the preparation of

8-chloro-2,3-benzodiazepine derivatives with AMPA/kainate receptor inhibiting activity

INVENTOR(S): Barkoczy, Jozsef; Ling, Istvan; Simig, Gyula; Szenasi,

Gabor; Gigler, Gabor; Kertesz, Szabolcs; Szuecs, Gyula; Szabo, Geza; Vegh, Miklos; Harsing, Laszlo

Gabor

PATENT ASSIGNEE(S): Egis Gyogyszergyar Rt, Hung.

SOURCE: PCT Int. Appl., 47 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

					KIND DATE				APPLICATION NO.					DATE					
	2005														2	0040	729		
	W: AE, AG,				AM,	ΑT,	ΑU,	ΑZ,	BA,	BB,	BG,	BR,	BW,	BY,	BZ,	CA,	CH,		
		CN,	CO,	CR,	CU,	CZ,	DE,	DK,	DM,	DZ,	EC,	EE,	EG,	ES,	FΙ,	GB,	GD,		
											JP,								
											MK,								
		NO,	NΖ,	OM,	PG,	PH,	PL,	PT,	RO,	RU,	SC,	SD,	SE,	SG,	SK,	SL,	SY,		
		ΤJ,	TM,	TN,	TR,	TT,	TZ,	UA,	UG,	US,	UZ,	VC,	VN,	YU,	ZA,	ZM,	ZW		
	RW:										SL,								
								•			BE,					-			
											LU,								
					BF,	ВJ,	CF,	CG,	CI,	CM,	GΑ,	GN,	GQ,	GW,	ML ,	MR,	ΝE,		
		- ,	TD,																
	2003		-		A2		2005			HU 2	003-	2449			20030804				
		2003002449												00040700					
		2004261491												20040729					
	2004261491									~- ^	004	0504	4 = 0	20040720					
	2534								CA 2004-2534458										
	1660				B1 20081119					EP 2004-769081					2	0040	729		
EP	1660	-						-	GB, GR, IT, LI, LU,						0.0		D		
	K:																	ш	
CNI	1071		51,	ш⊥,	ь∨, А						TR,							нк	
	1871 1005		_				20061129 CN 2004-80025409 20091021								20040729				
	2007						2009			JP 2006-522422					20040729				
	2007						2007					20040729							
	4146		70		Т		2007					20040729							
					됴		2000												
	PT 1660462 ES 2317038				T3		2009			PT 2004-769081					20040729				
KR 2006120578							2005	-		KR 2	004	7000	464	20040729 20060204					
	1094						2006			BG 2	006-	1094	62	20060302					
	2008													20080220					
	ZOOU (APP				111		_000	0020						A 20030804					
O-11-				• •							004-					0040			

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT OTHER SOURCE(S): CASREACT 142:219316; MARPAT 142:219316

GΙ

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The invention relates to new 8-chloro-2,3-benzodiazepine derivs. I [R = C1-4-alkyl (especially, Me or Et), NHR'; R' = C1-6-alkyl, C3-7-cycloalkyl] pharmaceutically acceptable acid addition salts thereof. The invention also encompasses a process for the preparation of said compds., pharmaceutical compns. containing them and new intermediates useful for the preparation of the new

8-chloro-2,3-benzodiazepine derivs. The said process comprises: (a) reducing nitrophenyl-2,3-benzodiazepine derivs. II; or (b) amidation of carboxylic acid derivs. III (Y = leaving group; Z = NH2, NO2) with NH2R'; (c) III (Z = NO2) is prepared from 8-chloro-2,3-benzodiazepine IV. Thus, amide I (R = NHMe) was prepared from 8-chloro-2,3-benzodiazepine via alkoxycarbonylation with ClCO2Ph, amidation with MeNH2, and reduction with Raney Ni. The compds. according to the invention possess AMPA/kainate receptor inhibiting activity. The bioactivity of I (R = NHMe) was determined [neuroprotective effect = -5 at 0.1 mg/kg i.p. against permanent focal cerebral ischemia in mice; ED50 = 2.5 μ M in spreading depression test in chicken retina; ED50 = 4.1 0.1 mg/kg i.p. in maximal electroshock test in mice; BWG = +9.9 g in toxicity test in rats].

IT 840526-71-8, 8-Chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-5H-2,3-benzodiazepine

RL: RCT (Reactant); RACT (Reactant or reagent)
(acylation and alkoxycarbonylation of; preparation of
8-chloro-2,3-benzodiazepine derivs. with AMPA/kainate receptor inhibiting activity)

RN 840526-71-8 CAPLUS

CN 5H-2,3-Benzodiazepine, 8-chloro-4-methyl-1-(3-methyl-4-nitrophenyl)- (CA INDEX NAME)

IT 840526-62-7P, 3-Acetyl-8-chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepine 840526-63-8P, 8-Chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-3-propionyl-3H-2,3-benzodiazepine

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and reduction of; preparation of 8-chloro-2, 3-benzodiazepine derivs.

with AMPA/kainate receptor inhibiting activity)

RN 840526-62-7 CAPLUS

CN Ethanone, 1-[8-chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

RN 840526-63-8 CAPLUS

CN 1-Propanone, 1-[8-chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

IT 840526-67-2P, 3-Acetyl-1-(4-amino-3-methylphenyl)-8-chloro-4-methyl-3H-2,3-benzodiazepine 840526-68-3P,

1-(4-Amino-3-methylphenyl)-8-chloro-4-methyl-3-propionyl-2,3-

benzodiazepine 840526-69-4P, 1-(4-Amino-3-methylphenyl)-8-chloro-4-methyl-3H-2,3-benzodiazepine-3-

carboxylic acid N-methylamide 840526-70-7P, 1-(4-Amino-3-methylphenyl)-8-chloro-4-methyl-3H-2,3-benzodiazepine-3-

carboxylic acid N-cyclopropylamide RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES

(preparation of 8-chloro-2,3-benzodiazepine derivs. with AMPA/kainate receptor inhibiting activity)

RN 840526-67-2 CAPLUS

(Uses)

CN Ethanone, 1-[1-(4-amino-3-methylphenyl)-8-chloro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

RN 840526-68-3 CAPLUS

CN 1-Propanone, 1-[1-(4-amino-3-methylphenyl)-8-chloro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

RN 840526-69-4 CAPLUS

CN 3H-2,3-Benzodiazepine-3-carboxamide, 1-(4-amino-3-methylphenyl)-8-chloro-N,4-dimethyl- (CA INDEX NAME)

RN 840526-70-7 CAPLUS

CN 3H-2,3-Benzodiazepine-3-carboxamide, 1-(4-amino-3-methylphenyl)-8-chloro-N-cyclopropyl-4-methyl- (CA INDEX NAME)

ΙT 840526-64-9P, 8-Chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepine-3-carboxylic acid phenyl ester 840526-65-0P , 8-Chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepine-3-carboxylic acid N-methylamide 840526-66-1P, 8-Chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepine-3carboxylic acid N-cyclopropylamide RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (preparation of 8-chloro-2,3-benzodiazepine derivs. with AMPA/kainate receptor inhibiting activity) 840526-64-9 CAPLUS RN CN 3H-2,3-Benzodiazepine-3-carboxylic acid, 8-chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-, phenyl ester (CA INDEX NAME)

RN 840526-65-0 CAPLUS
CN 3H-2,3-Benzodiazepine-3-carboxamide,
8-chloro-N,4-dimethyl-1-(3-methyl-4-nitrophenyl)- (CA INDEX NAME)

RN 840526-66-1 CAPLUS
CN 3H-2,3-Benzodiazepine-3-carboxamide,
8-chloro-N-cyclopropyl-4-methyl-1-(3-methyl-4-nitrophenyl)- (CA INDEX NAME)

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2011 ACS on STN

ACCESSION NUMBER: 2002:487537 CAPLUS

DOCUMENT NUMBER: 137:63266

TITLE: Preparation of 2,3-benzodiazepines as AMPA

antagonists.

INVENTOR(S): Ling, Istvan; Barkoczy, Jozsef; Simig, Gyula; Greff,

Zoltan; Ratkai, Zoltan; Szabo, Geza; Vegh, Miklos; Gigler, Gabor; Szenasi, Gabor; Martonne Marko, Bernadett; Levay, Gyoergy; Harsing, Laszlo Gabor

PATENT ASSIGNEE(S): Egis Gyogyszergyar Rt., Hung.

SOURCE: PCT Int. Appl., 40 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PA:	FENT 1	NO.			KIND DATE			APPLICATION NO.						DATE				
					A1 20020627													
W: AE, AG,			AL,	AM,	AT,	AU,	AZ,	BA,	BE	B, BG,	BR,	BY,	BZ,	CA,	CH,	CN,		
		CO,	CR,	CU,	CZ,	DE,	DK,	DM,	DZ,	EC	C, EE,	ES,	FΙ,	GB,	GD,	GE,	GH,	
		GM,	HR,	ID,	IL,	IN,	IS,	JP,	ΚE,	KG	KP,	KR,	KΖ,	LC,	LK,	LR,	LS,	
		LT,	LU,	LV,	MA,	MD,	MG,	MK,	MN,	MW	I, MX,	ΜZ,	NO,	NZ,	OM,	PH,	PL,	
		PT,	RO,	RU,	SD,	SE,	SG,	SI,	SK,	SI	, TJ,	TM,	TN,	TR,	ΤT,	TZ,	UA,	
		,				,	ZA,											
	RW:										I, TZ,							
											I, IT,							
											Q, GW,							
HU	2000	0049	94		A2		2002	1128		HU	2000-		20001221					
	2251																	
CA	2431		A1		2002	0627	CA 2001-2431761						20011219					
CA	A 2431761				С	2010	0525	0000 15056						00011010				
AU	AU 2002017356				А		2002	0701	AU 2002-17356 EP 2001-271364						20011219			
EP	1351	942			A1	2003	1015	EP 2001-271364						2	0011	219		
		942			В1		2004	0721										
	R:										R, IT,	LI,	LU,	NL,	SE,	MC,	PT,	
		IE,	SI,	LT,	LV,	FI,	RO,	MK,	CY,	AL	TR							
JP	2004	5162	84		T		2004	0603		JP	2002-	5515	41		2	0011	219	
JP	4201	252			B2		2008	1224		3 m	2002-	0710	C 1		^	0011	010	
AT	2/15	48			T		2004	0815		AT	2001-	.2/13	64		2	0011	219	
PT	1321	942			E		2004	1130		PT	2001-	2713	64		2	0011	219	
	2225										2001-					0011		
	1212 2866						2007	0228		KU CE	2003- 2003-	700			2	0011		
		∠ U 7 1			D6		2009	0203		OZ VC	2003-	1600			2	0011	219	
CZ 301371 BG 107932					7/		2010	0203	CZ 2003-1680 BG 2003-107932						20030630			
) <u>/</u> n n a ?	510		A. 7\1		2004	044 <i>1</i> 0513		DG IIC	2003-	.1500	92 98		20030620			
0.0	7189	0094. 711	210		B2		2004	U313		U.S	2003-	.4000	20		4	0031	04/	
	/109 Y APP:				שט		2007	0010		пп	2000-	./199/			Δ 2	0001	221	
/1\	L AFF.	□1/1 •	TMEO	• •							2000-					0001		
										MO	2001-	11010	_		v	O O T T	41	

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 137:63266

GI

AB Title compds. (I; X = H, Cl, MeO; Y = H, halo; Z = Me, Cl; R = alkyl, NR1R2; R1, R2 = H, alkyl, alkoxy, cycloalkyl), were prepared Thus, 3-acetyl-4,5-dihydro-8-chloro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepine in methanol/CH2Cl2 was stirred with wet Raney nickel and hydrazine hydrate for 45 min to give 49% 3-acetyl-1-(4-amino-3-methylphenyl)-4,5-dihydro-8-chloro-4-methyl-3H-2,3-benzodiazepine. The latter prolonged the survival time of MgCl2-treated mice with PD50 = 4.6 mg/kg i.p.

IT 439143-67-6P 439143-68-7P 439143-70-1P 439143-71-2P

Ι

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of 2,3-benzodiazepines as AMPA antagonists)

RN 439143-67-6 CAPLUS

CN Ethanone, 1-[1-(4-amino-3-methylphenyl)-8-chloro-4,5-dihydro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

RN 439143-68-7 CAPLUS

CN 1-Propanone, 1-[1-(4-amino-3-methylphenyl)-8-chloro-4,5-dihydro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

RN 439143-70-1 CAPLUS

CN Ethanone, 1-[1-(4-amino-3-methylphenyl)-7,8-dichloro-4,5-dihydro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

RN 439143-71-2 CAPLUS

CN 1-Propanone, 1-[1-(4-amino-3-methylphenyl)-7,8-dichloro-4,5-dihydro-4-methyl-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

IT 439143-77-8 439143-78-9 439143-80-3 439143-81-4

RL: RCT (Reactant); RACT (Reactant or reagent) (preparation of 2,3-benzodiazepines as AMPA antagonists)

RN 439143-77-8 CAPLUS

CN Ethanone, 1-[8-chloro-4,5-dihydro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

RN

439143-78-9 CAPLUS 1-Propanone, 1-[8-chloro-4,5-dihydro-4-methyl-1-(3-methyl-4-nitrophenyl)-CN 3H-2,3-benzodiazepin-3-y1]- (CA INDEX NAME)

439143-80-3 CAPLUS RN

Ethanone, 1-[7,8-dichloro-4,5-dihydro-4-methyl-1-(3-methyl-4-nitrophenyl)-1-(3-methyl-4-nitrophenyl)CN 3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

10/567,598

RN 439143-81-4 CAPLUS CN 1-Propanone, 1-[7,8-dichloro-4,5-dihydro-4-methyl-1-(3-methyl-4-nitrophenyl)-3H-2,3-benzodiazepin-3-yl]- (CA INDEX NAME)

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT